arXiv:1105.1810v1 [math.AP] 9 May 2011

A Liouville comparison principle for entire sub- and super-solutions of the equation $u_t - \Delta_p(u) = |u|^{q-1}u$.

Vasilii V. Kurta.

January 20, 2013

Abstract

We establish a Liouville comparison principle for entire sub- and super-solutions of the equation $(*)\ w_t - \Delta_p(w) = |w|^{q-1}w$ in the half-space $\mathbb{S} = \mathbb{R}^1_+ \times \mathbb{R}^n$, where $n \geq 1$, $q > 0$ and $\Delta_p(w) := \mathrm{div}_x \left(|\nabla_x w|^{p-2}\nabla_x w\right)$, $1 < p \leq 2$. In our study we impose neither restrictions on the behaviour of entire sub- and super-solutions on the hyper-plane $t = 0$, nor any growth conditions on the behavior of them or any of their partial derivatives at infinity. We prove that if $1 < q \leq p - 1 + \frac{p}{n}$, and u and v are, respectively, an entire weak super- and an entire weak sub-solution of $(*)$ in $\mathbb{S}$ which belong, only locally in $\mathbb{S}$, to the corresponding Sobolev space and are such that $u \leq v$, then $u \equiv v$. The result is sharp. As direct corollaries we obtain both new and known Fujita-type and Liouville-type results.

1 Introduction and Definitions.

The purpose of this work is to obtain a Liouville comparison principle of elliptic type for entire weak sub- and super-solutions of the equation

$$w_t - \Delta_p(w) = |w|^{q-1}w \tag{1}$$

in the half-space $\mathbb{S} = (0, +\infty) \times \mathbb{R}^n$, where $n \geq 1$ is a natural number, $q > 0$ is a real number and $\Delta_p(w) := \mathrm{div}_x\left(|\nabla_x w|^{p-2}\nabla_x w\right)$, $1 < p \leq 2$, defines the well-known p-Laplacian operator. Under entire sub- and super-solutions of (1) we understand sub- and super-solutions of (1) defined in the whole half-space $\mathbb{S}$ and under Liouville theorems of elliptic type we understand Liouville-type theorems which, in their formulations, have no restrictions on the behaviour of sub- or super-solutions to the parabolic equation (1) on the hyper-plane $t = 0$. We would also like to underline that we impose no growth conditions on the behavior of sub- or super-solutions of (1), as well as of all their partial derivatives, at infinity.

Definition 1 *Let $n \geq 1$, $p > 1$ and $q > 0$. A function $u = u(t, x)$ defined and measurable in $\mathbb{S}$ is called an entire weak super-solution of the equation (1) in $\mathbb{S}$ if it belongs to the function space $L_{q,loc}(\mathbb{S})$, with $u_t \in L_{1,loc}(\mathbb{S})$ and $|\nabla_x u|^p \in L_{1,loc}(\mathbb{S})$, and satisfies the integral inequality*

$$\int_{\mathbb{S}} \left[u_t \varphi + \sum_{i=1}^{n} |\nabla_x u|^{p-2} u_{x_i} \varphi_{x_i} - |u|^{q-1} u \varphi \right] dt dx \geq 0 \tag{2}$$

for every non-negative function $\varphi \in C^\infty(\mathbb{S})$ with compact support in $\mathbb{S}$, where $C^\infty(\mathbb{S})$ is the space of all functions defined and infinitely differentiable in $\mathbb{S}$.

Definition 2 *A function $v = v(t, x)$ is an entire weak sub-solution of (1) if $u = -v$ is an entire weak super-solution of (1).*

2 Results.

Theorem 1 *Let $n \geq 1$, $2 \geq p > 1$ and $1 < q \leq p - 1 + \frac{p}{n}$, and let u be an entire weak super-solution and v an entire weak sub-solution of (1) in $\mathbb{S}$ such that $u \geq v$. Then $u \equiv v$ in $\mathbb{S}$.*

The result in Theorem 1, which evidently has a comparison principle character, we term a Liouville-type comparison principle, since, in the particular cases when $u \equiv 0$ or $v \equiv 0$, it becomes a Liouville-type theorem of elliptic type, respectively, for entire sub- or super-solutions of (1).

Since in Theorem 1 we impose no conditions on the behaviour of entire sub- or super-solutions of the equation (1) on the hyper-plane $t = 0$, we can formulate, as a direct corollary of the result in Theorem 1, the following comparison principle, which in turn one can term a Fujita comparison principle, for entire sub- and super-solutions of the Cauchy problem for the equation (1). It is clear that in the particular cases when $u \equiv 0$ or $v \equiv 0$, it becomes a Fujita-type theorem, respectively, for entire sub- or super-solutions of the Cauchy problem for the equation (1).

Theorem 2 *Let $n \geq 1$, $2 \geq p > 1$ and $1 < q \leq p - 1 + \frac{p}{n}$, and let u be an entire weak super-solution and v an entire weak sub-solution of the Cauchy problem, with possibly different initial data for u and v, for the equation (1) in $\mathbb{S}$ such that $u \geq v$. Then $u \equiv v$ in $\mathbb{S}$.*

Note that the results in Theorems 1 and 2 are sharp. The sharpness of these for $q > p - 1 + \frac{p}{n} \geq 1$ follows, for example, from the existence of non-negative self-similar entire solutions to (1) in $\mathbb{S}$, that was shown in [1]. Also, there one can find a Fujita-type theorem on blow-up of non-negative entire solutions of the Cauchy problem for (1), which was obtained as a very interesting generalization of the famous blow-up result in [2] to quasilinear parabolic equations. For $0 < q \leq 1$, it is evident that the function $u(t,x) = e^t$ is a positive entire classical super-solution of (1) in $\mathbb{S}$.

To prove the results in Theorems 1 and 2 we further develop the approach that was proposed for solving similar problems for semilinear parabolic equations in [3]. A new key point in our proof is using the fact that for $1 < p \leq 2$ the p-Laplacian operator Δ_p satisfies the α-monotonicity condition (see, e.g., [4]) with $\alpha = p$.

Acknowledgments.

This research is financially supported by the Alexander von Humboldt Foundation (AvH). The author is very grateful to AvH for the opportunity to visit the Mathematical Institute of Köln University and to Professor B. Kawohl for his cordial hospitality during this visit.

References

[1] V.A. Galaktionov and H.A. Levine, *A general approach to critical Fujita exponents in nonlinear parabolic problems*, Nonlinear Anal. (1998) v. 34, no. 7, 1005–1027.

[2] H. Fujita, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$*, J. Fac. Sci. Univ. Tokyo, Sect. I (1966), v. 13, 109–124.

[3] A.G. Kartsatos and V.V. Kurta, *On a comparison principle and the critical Fujita exponents for solutions of semilinear parabolic inequalities*, J. London Math. Soc. (2) 66 (2002), no. 2, 351–360.

[4] V.V. Kurta, *Comparison principle for solutions of parabolic inequalities*, C. R. Acad. Sci. Paris, Série I, 322 (1996), 1175–1180.

Author's address:

Vasilii V. Kurta
Mathematical Reviews
416 Fourth Street, P.O. Box 8604
Ann Arbor, Michigan 48107-8604, USA
e-mail: vkurta@umich.edu, vvk@ams.org